

HOX Homeobox Genes Exhibit Spatial and Temporal Changes in Expression During Human Skin Development

Eric J. Stelnicki,* László G. Kömüves,†** Angela O. Kwong,†** Dennis Holmes,‡ Peter Klein,§ Sophia Rozenfeld,§** H. Jeffrey Lawrence,§** N. Scott Adzick,¶ Michael Harrison,* and Corey Largman§**

Departments of *Surgery, †Dermatology, and §Medicine, University of California, San Francisco, California, U.S.A.; ‡Department of Surgery, Washington University Medical Center, St Louis, Missouri, U.S.A.; ¶Children's Hospital and University of Pennsylvania, Philadelphia, Pennsylvania, U.S.A.; and

**Departments of Medicine and Dermatology, San Francisco VA Medical Center, San Francisco, California, U.S.A.

The spatial and temporal deployment of *HOX* homeobox genes along the spinal axis and in limb buds during fetal development is a key program in embryonic pattern formation. Although we have previously reported that several of the *HOX* homeobox genes are expressed during murine skin development, there is no information about developmental expression of *HOX* genes in human skin. We have now used reverse transcriptase polymerase chain reaction, in conjunction with a set of degenerate oligonucleotide primers, to identify a subset of *HOX* genes that are expressed during human fetal skin development. *In situ* hybridization analyses demonstrated that there were temporal and spatial shifts in expression of these genes. Strong *HOXA4* expression was detected in the

basal cell layers of 10 wk fetal epidermis and throughout the epidermis and dermis of 17 wk skin, whereas weak signal was present in the granular layer of newborn and adult skin. The expression patterns of *HOXA5* and *HOXA7* were similar, but their expression was weaker. *In situ* hybridization analysis also revealed strong *HOXC4* and weaker *HOXB7* expression throughout fetal development, whereas *HOXB4* was expressed at barely detectable levels. Differential *HOX* gene expression was also observed in developing hair follicles, and sebaceous and sweat glands. None of the *HOX* genes examined were detected in the adult dermis. **Key words:** dermis/epidermis/gene regulation/homeodomain. *J Invest Dermatol* 110:110–115, 1998

Human skin is comprised of a self-renewing epidermis derived from ectodermal tissue, and an underlying dermis, which is of mesodermal origin. There is increasing evidence that BMP-4 (Nohno *et al*, 1995a), and other signaling molecules such as sonic hedgehog (Nohno *et al*, 1995b), TGF- α (Luetke *et al*, 1993; Mann *et al*, 1993), and EGF (Miettinen *et al*, 1995; Murillas *et al*, 1995), play important inductive roles in epithelial-mesenchymal signaling during skin development and hair formation. It has been hypothesized that a set of transcription factors must mediate the effects of these putative morphogens to regulate skin-specific gene transcription. In this regard, the Lef-1 transcription factor has been shown to play a role in whisker and hair formation (van Genderen *et al*, 1994; Zhou *et al*, 1995); however, the molecular mechanisms that regulate skin and hair development remain largely unknown.

The *HOX* homeobox genes appear to be candidates for the regulation of skin development. These master developmental regulatory genes encode proteins that contain a conserved 60 amino acid, DNA binding homeodomain and function as transcription factors (Gehring *et al*, 1994). The 39 human homeobox genes are arrayed in four parallel chromosomal clusters such that individual genes from each loci (e.g., *HOXA4*, *HOXB4*, *HOXC4*, and *HOXD4*) show a high degree of homology within their putative homeodomains. The *HOX* homeo-

domain proteins appear to regulate the formation of multiple structures along various body axes such as the spinal cord, and limbs (Krumlauf, 1994). We (Detmer *et al*, 1993; Mathews *et al*, 1993) and others (Bieberich *et al*, 1991; Reiger *et al*, 1994) have previously detected expression of several of the *Hox* genes, particularly those in the *Hox-b* cluster, in fetal and adult murine skin. In addition, there have been several reports of spatially restricted expression of *Hox* genes in developing murine (Bieberich *et al*, 1991; Kanzler *et al*, 1994) and chicken skin (Chuong *et al*, 1990). Based on these studies and the well-recognized role of homeodomain proteins in development, Scott and Goldsmith proposed that the *HOX* homeobox genes play a major regulatory role in human skin development (Scott and Goldsmith, 1993). Until now, however, there have not been studies on the spatial and temporal expression of *HOX* genes in developing human skin. We report here the use of a reverse transcriptase polymerase chain reaction (RT-PCR) based approach to identify which of the 39 human *HOX* genes are expressed in first and second trimester human fetal skin and which are expressed in adult skin. Only a small subset of *HOX* genes were detected using this method. *In situ* hybridization analysis revealed spatial and temporal changes in the expression of these genes within both the developing epidermis and the dermis, as well as in hair follicles.

MATERIALS AND METHODS

Tissue samples Tissue samples were obtained under NIH guidelines and after approval of the University of California San Francisco Committee on Human Research. Tissue was fixed in 4% paraformaldehyde for 4 h, dehydrated in graded ethanol, and embedded in paraffin. For some tissues, half was frozen in solution D (see below) for subsequent RNA isolation. Following informed consent newborn and adult skin samples were obtained from the margins of specimens from elective surgery.

Manuscript received September 16, 1996; revised August 20, 1997; accepted for publication October 14, 1997.

Reprint requests to: Dr. Corey Largman (151H), VA Medical Center, 4150 Clement Street, San Francisco, CA 94121.

Abbreviation: RT-PCR, reverse transcriptase polymerase chain reaction.

Dermo-epidermal separation Five samples of second trimester fetal scalp skin (18–22 wk) and five samples of adult skin (50–65 y) were subjected to dermal/epidermal separation by exposure of full thickness skin samples to dry heat (60°C for 60 s) (Holleran *et al.*, 1992). The epithelium was then scraped off the dermis and placed in a separate container for RNA isolation.

RNA isolation Total RNA was isolated from fetal, neonatal, and adult skin using the guanidinium thiocyanate method (Chomczynski *et al.*, 1987). Skin from the scalp, back, or limbs was homogenized in solution D (4 M guanidinium thiocyanate, 25 mM sodium citrate, pH 7, 0.5% sarcosyl, 0.1% 2-mercaptoethanol), and RNA was separated from DNA and protein by extraction with acid phenol:chloroform:isoamyl alcohol (50:49:1). RNA concentrations were estimated by the optical density at 260 nm, and RNA was stored in 70% ethanol at –70°C.

RT-PCR RT-PCR was used to survey for *HOX* gene expression in each sample. We used a protocol that included two polymerase chain reaction (PCR) amplifications (Savageau *et al.*, 1994). First, amplified cDNA representing total mRNA was synthesized from each tissue RNA. A subsequent PCR amplification using degenerate primers designed against all 39¹ *HOX* genes permitted their identification by cloning and sequencing of the PCR reaction products. The first strand cDNA was synthesized by adding total RNA (1 µg) to 10×reverse transcriptase buffer, dNTP, Superscript II reverse transcriptase (Gibco BRL, Gaithersburg, MD), and an oligo dT primer: (5′-CATGTCGTCAGGCCG-CTCTGGACAAAATATGAATTC (T)₂₁-3′). Following reverse transcription and phenol/chloroform extraction, a 5′ polyA tail was added to the cDNA using terminal transferase. The same oligo dT primer was then re-added to the solution (1 µg per µl) and PCR was performed (94°C, 2 min; 55°C, 3 min; 72°C, 10 min) for 35 cycles.

The amplified cDNA mixtures were screened using a set of degenerate *HOX* gene primers derived from two conserved regions of the homeobox, separated by 118 nt: (5′-GA (G/A) (T/C)T (G/A/T)GA (A/G) (C/A/G) (G/A) (G/A/C)GA (A/G)T (A/T) (T/C)-3′) and (5′-NC (G/T) (A/G)TT (T/C)TG (A/G)AACCANA (C/T)-3′). A Hot Wax PCR kit (10 × buffer pH = 10.0 and 1.5 MgCl beads) (Invitrogen, La Jolla, CA) was used to maximize the PCR reaction (94°C, 2 min; 48°C, 2.5 min; and 72°C, 3 min) for 32 cycles. This procedure produced a 118 bp band that was detected on a 2% agarose gel (FMC, Rockland, ME) and that contained amplified products reflecting the homeobox gene transcripts present in each tissue sample.

In order to identify the various homeobox cDNA derived from each sample, the 118 bp DNA fragments were cut from the gels, purified (Bio 101, Vista, CA), and ligated into the T/A PcrII vector (Invitrogen). Insert-containing clones were sequenced using ³⁵S-ATP and the Sanger dideoxy method (Promega, Madison, WI). Sequences were identified by comparison to the known 38 human *HOX* genes (Genbank) using GCG-8 software (Genetics Computer Group, Madison, WI).

RNA template labeling, probe preparation, and *in situ* hybridization Templates for human *HOXA4*, *HOXA5*, *HOXA7*, *HOXB4*, *HOXB7*, and *HOXC4* were constructed by subcloning 3′ nonhomeobox fragments in Bluescript SK⁺ (Stratagene, La Jolla, CA). Linearized plasmids (1 µg) were used to synthesize digoxigenin-labeled sense and anti-sense RNA probes (Boehringer, Indianapolis, IN). RNA *in situ* hybridization was performed as described (Mathews *et al.*, 1993) with modifications to use digoxigenin labeled-probes (Wilkinson, 1992). Sections were deparaffinized in three changes of Americlear (Baxter Diagnostics, Deerfield, IL) and re-hydrated through graded ethanol. Following washing, sections were digested with proteinase K (10 µg per ml) for 30 min at 37°C, rinsed in physiologically buffered saline, washed in 0.1 M triethanolamine with 0.25% acetic anhydride, prehybridized in 50% formamide, 2 × sodium citrate/chloride buffer for 1 h at 37°C, and then allowed to air dry. The digoxigenin-labeled probes were diluted in hybridization solution (2 × sodium citrate/chloride buffer, 12.5 × Denhardt's solution, 50% formamide, 0.5% sodium dodecyl sulfate, 0.25 µg salmon sperm per ml, 0.5% sodium pyrophosphate, 10 mM TrisHCl, pH 7.4), applied to each section, which was then covered with parafilm and incubated overnight at 42°C. Sections were washed, treated with RNase A (20 µg per ml), washed in a series of increasing




Figure 1. *HOX* homeobox genes are expressed in developing skin. (A) PCR analysis, using degenerate primers designed to amplify a region of the homeobox from any of the known 38 human *HOX* genes, was used to detect *HOX* transcripts in whole second trimester fetal skin. Following an initial RT-PCR amplification using oligo dT primers to generate an amplified cDNA, a second round of PCR was performed with the degenerate primers to generate a 118 bp DNA fragment (→, lane 2) reflecting the presence of *HOX* transcripts. Lane 1 is an identical sample in which reverse transcriptase was omitted from the initial cDNA synthesis step. (B) Identification of *HOX* gene transcripts in whole skin samples. The 118 bp PCR products from each skin sample, prepared as shown in (A), were cloned and sequenced, and comparison with published sequences permitted identification of *HOX* transcripts. The values shown represent the number of clones that matched a given *HOX* gene sequence. Expression of the other 27 known human *HOX* homeobox genes was not detected. (C) *HOX* genes were not detected in adult dermis. The degenerate primer PCR method was also used to amplify *HOX* transcripts from fetal and adult skin samples that have been split into epidermis and dermis. Appropriately sized PCR products were detected in both fetal and adult epidermis (→, lanes 3 and 4), as well as in fetal dermis (lane 1); however, a 118 bp band was not detected in the adult dermis (lane 2). Cloning and sequencing of the smeared PCR products from the adult dermis confirmed that no *HOX* transcripts could be detected. In contrast, the PCR bands for the fetal and adult epidermis and the fetal dermis all gave data similar to those reported in (B).

stringency solutions, and rinsed in buffer (100 mM TrisHCl, pH 7.5, 150 mM NaCl). Sections were incubated (1 h) in anti-digoxigenin-horseradish peroxidase antibody (1:500) in buffer [1% non-fat dried milk, 0.5% fish skin gelatin, 10 mM 500 mM NaCl, 0.1% Tween 20, TrisHCl (pH 7.6)], then incubated with biotinylated-tyramide (15 min), followed by a streptavidin-peroxidase (15 min) (Dako, Carpinteria, CA). Signals were visualized with DAB chromogen substrate (5 min). For some *HOXA5* experiments, the color development time was 25 min. To detect *HOXA7* signals, color development was for 25 min and a DAB color enhancer (Innovex Biosciences, Richmond, CA) was added to the reaction. Some sections were counter-stained with hematoxylin to visualize tissue morphology. Photographs were taken with a Nikon Microphot FX photomicroscope and digitized using a Nikon Coolscan slide scanner. Composite images were created with Adobe Photoshop, and printed on a Kodak dye sublimation printer. Except for background color adjustment, no other image processing was performed.

RESULTS

A subset of *HOX* homeobox genes is expressed in developing human fetal skin We first used an RT-PCR/sequence analysis to identify *HOX* homeobox genes expressed in human fetal and adult scalp skin. With this method, an initial RT-PCR-amplified cDNA library representing the total mRNA of each tissue sample was constructed. This cDNA was then subjected to a second round of PCR amplification using a set of degenerate PCR primers, derived from conserved portions of the homeobox region, which were designed to amplify all of the human *HOX* genes. **Figure 1A** shows a representative experiment in which the expected 118 bp PCR product band was detected in RNA isolated from whole second trimester fetal scalp skin. The 118 bp PCR product band derived from each skin sample was excised from the gel, cloned, and sequenced. Clones containing *HOX* gene transcripts were identified by comparison with published sequences of the 38 known *HOX* genes.¹ When the PCR products were cloned and identified, the majority of the *HOX* gene transcripts detected in fetal skin (**Fig 1B**) belonged to the *HOX A*

¹At the start of this project, 38 *HOX* genes had been described. As part of this work, an additional human *HOX* gene was identified in developing dermis. The new gene, *HOXB13*, appears to be differentially regulated in scarless fetal wound healing and is the subject of a submitted manuscript (Stelnicki EJ, Arbeit J, Cass DL, Saner C, Harrison M, Largman C: Changes in expression of the human homeobox genes PRX-2 and *HOXB13* are associated with scarless fetal wound healing. Manuscript submitted). This gene was independently isolated by Zeltser *et al.* (1996) during these studies. The degenerate primers used contained a match for the *HOXB13* homeobox.

cluster. Of these, *HOXA4*, *HOXA5*, and *HOXA7* appeared to be predominant. Transcripts for *HOXA6*, *HOXA10*, *HOXA11*, and *HOXB7* were also detected with lower frequency in whole fetal skin. It must be emphasized that it is not possible to use these data to compare the relative levels of expression between time points. Like the fetal tissue, the majority of *HOX* gene transcripts detected in the human neonatal scalp were *HOXA4*, *HOXA5*, and *HOXA7* (Fig 1B). This same trend continued in the adult tissue, with the additional detection of several members of the *HOX-B* cluster (*HOXB1*, *HOXB3*, and *HOXB6*) and one member of the *HOX-C* locus (*HOXC4*) (Fig 1B). The RT-PCR analysis was repeated on RNA isolated from back and arm skin from first and second trimester embryos as well as from newborn foreskin. These experiments revealed the same pattern in which *HOXA4*, *HOXA5*, and *HOXA7* were the major transcripts detected (data not shown).

HOX gene transcripts are not detected in adult dermis by RT-PCR A similar RT-PCR analysis was performed on fetal and adult skin samples that were split into epidermis and dermis. Although the expected 118 bp PCR band representing amplification of *HOX* transcripts was detected in fetal epidermis and dermis, as well as in adult epidermis, there was no apparent *HOX* gene expression in adult dermis as reflected by the absence of the appropriate size PCR product in multiple ($N = 5$) samples (Fig 1C, lane 2). When the PCR bands from the fetal epidermis and dermis and the adult epidermis were cloned and sequenced, data similar to those shown in Fig 1B were obtained (not shown). As anticipated by the lack of a 118 bp RT-PCR band, cloning of the PCR products from adult dermis did not yield identifiable *HOX* gene sequences.

HOXA4 is expressed in a temporal and spatial pattern during epidermal development In order to obtain spatial localization of *HOX* gene transcripts, we used *in situ* hybridization to analyze expression during skin development. We analyzed *HOX* gene expression in skin from the scalp, back, and arm from first trimester (10 wk estimated gestational age), early second trimester (14 wk estimated gestational age), mid-second trimester (17 wk estimated gestational age), and late second trimester (21 wk estimated gestational age) embryos, as well as from normal newborn (4 mo old) and adult skin samples. Because *HOXA4*, *HOXA5*, and *HOXA7* appeared from the RT-PCR data to be strongly expressed in developing skin, we first examined expression of these genes at each developmental time point (Fig 2). Because the signals for *HOXA4* were quite strong, these sections were counter-stained with hematoxylin to visualize tissue morphology (Fig 2A–F). Weaker *HOXA5* (Fig 2G–L) and *HOXA7* (Fig 2M–R) signals were visualized on adjacent tissue sections without counter-staining. In order to be able to compare *HOXA4* expression with that of the other genes, a subset of sections was also examined for *HOXA4* signals in the absence of counter-staining (Fig 3A–C). *HOXA4* was predominantly expressed in the basal cell layer, and was localized to the lower layers of the developing epidermis in the early (10 wk) embryonic skin (Figs 2A, 3A). Control slides in which tissues were hybridized with a sense RNA probe were negative (see Fig 3R). Cells of the periderm (present at 10 and 14 wk) did not express *HOXA4*, or any other *HOX* gene studied. *HOXA4* expression appeared to gradually spread throughout the epidermis from 10 to 17 wk of development until by mid-second trimester, the signal was detected in the entire epidermis, except the stratum corneum (Figs 2B,C, 3B). It should be noted that the stratum corneum was uniformly negative at each developmental stage for every probe studied. By late second trimester *HOXA4* expression was less prevalent in the basal cell layer but was still detected throughout the suprabasal epidermis (Figs 2D, 3C). *HOXA4* expression was downregulated further in the basal and stratum spinosum layers of the newborn (Fig 2E) and adult skin (Fig 2F), where expression appeared to be localized to the upper granular layers of the epidermis. The specimens shown in Fig 2(A–F) were all from scalp except the 14 wk sample (Fig 2B) that was from the back. *In situ* hybridization analysis of *HOXA4* expression in the corresponding series of back or arm skin samples showed essentially identical temporal and spatial changes throughout development (data not shown). In addition, the relative expression levels of *HOXA4*, as

well as those of the other *HOX* genes described below, did not vary substantially between these three tissues during development.


HOXA4 expression is temporally regulated in the dermis during skin development *HOXA4* expression appeared to be virtually absent in the dermis of first trimester (week 10, Figs 2A, 3A) and early second trimester (week 14, Fig 2B) fetal skin; however, by mid-second trimester (week 17, Figs 2C, 3B) *HOXA4* expression was seen in dermal fibroblasts. Weak expression persisted at late second trimester dermis (week 21, Figs 2D, 3C), but was absent in newborn (Fig 2E) and adult (Fig 2F) dermis. Fibroblasts in the dermal papillae did not express *HOXA4* or other *HOX* genes studied (Figs 2B–D, 3M, see also the section on hair follicle development below).

HOXA5 and HOXA7 are expressed in similar patterns but at lower levels than HOXA4 in developing skin *In situ* hybridization experiments revealed that *HOXA5* (Fig 2G–L) was detected at lower levels but in a similar pattern to *HOXA4*, whereas initial experiments with a *HOXA7* probe gave very weak signals. We used increased color development time and an additional amplification step in order to localize expression of the *HOXA7* gene (Fig 2M–R), which appeared to follow similar temporal and spatial patterns to those observed for *HOXA4*. Thus, expression appeared to move from the basal layers in early epidermal development towards the outer epidermis in newborn and adult skin. Weak expression of *HOXA5* and *HOXA7* was detected in the fetal dermis during the second trimester of fetal development (Fig 2I–O), but no expression of these genes was observed in newborn (Fig 2K,Q) or adult dermis (Fig 2L,R).

HOXB4, HOXB7, and HOXC4 are variably expressed in developing skin We also used *in situ* hybridization to assess the expression patterns of *HOX* genes detected at lower levels by the RT-PCR method (*HOXB7* and *HOXC4*), as well as a gene not detected by this procedure (*HOXB4*). *HOXC4* was expressed in similar temporal and spatial patterns to those observed for *HOXA4* (compare Fig 3J–L with Fig 3A–C), whereas *HOXB4* was barely detectable (compare Fig 3D–F with 3R). *HOXB7* was weakly expressed in the basal and suprabasal layers of the 10 wk old skin (Fig 3G), was downregulated in early second trimester skin (Fig 3H), and appeared to be upregulated throughout the epidermis in 21 wk skin (Fig 3I). Expression of each of these genes was downregulated and restricted to the upper epidermis in newborn and adult skin (data not shown). Essentially identical results were obtained using skin from either the back or the arm (not shown).

Differential HOX gene expression during hair follicle, sweat gland, and sebaceous gland development As seen in Fig 2B, early hair follicle formation was observed in the 14 wk skin. At this time, none of the *HOX* genes studied could be detected in the prospective bud (Fig 2B,H,N, and data not shown); however, by 17 wk, clear expression of *HOXA4* (Figs 2C, 3B) and *HOXA5* (Fig 2I) was detected in developing follicles, whereas *HOXC4* expression was weak but noticeable (Fig 3K). None of the other *HOX* genes examined were detected in the developing hair follicles of the 17 wk skin. By 21 wk, signals for *HOXB7* in the maturing hair follicles (Fig 3I) were approximately equal to those observed for *HOXA4* (Figs 2D, 3C, and 3M) or *HOXC4* (Figs 3L and 3Q), whereas expression of the other genes was not detected. In newborn skin, the mature follicles showed expression of *HOXA4* (Figs 3M and 3O), *HOXA5* (Fig 3P), and *HOXA7* (Fig 2Q). More importantly, these *HOX* genes appeared to be expressed in somewhat different spatial patterns in the mature hair follicles in 21 wk and 4 mo skin. Thus *HOXA4* (compare Figs 3M and 3N, and see Fig 3O), *HOXA7* (Fig 2Q), and *HOXC4* (Fig 3Q) were expressed in all the epidermal layers of the hair follicle but not in the melanocytes or in the dermal papillae. In contrast, *HOXA5* expression was restricted to the inner root sheath (Fig 3P). Only extremely weak *HOX* gene expression was detected in the hair follicles within the adult skin (data not shown). In addition to prevalent expression in hair follicles, *HOXA4* was clearly also expressed in sweat glands in newborn skin (Fig 3O) and adult skin (data not shown). Expression of the other *HOX* genes studied was not detected in sweat glands at any stage of development. *HOXA4* was also clearly detected

Figure 2. *HOXA4*, *HOXA5*, and *HOXA7* genes are expressed in spatial and temporal patterns during skin development. *In situ* hybridization analysis was used to detect expression of *HOXA4* (A–F), *HOXA5* (G–L), and *HOXA7* (M–R) in serial sections from 10 wk scalp (A, G, and M), 14 wk back (B, H, and N), 17 wk scalp (C, I, and O), 21 wk scalp (D, J, and P), newborn scalp (E, K, and Q), and adult scalp (F, L, and R). Parts (A)–(F) were counterstained with hematoxylin to visualize tissue morphology. Parts (G)–(L) and (M)–(Q) represent adjacent sections to those stained in (A)–(F). Digoxigenin-labeled RNA probes were hybridized with tissue sections, and a tyramide amplification and peroxidase detection was used to visualize *HOX* gene expression as described in *Materials and Methods*. Whereas detection of *HOXA4* was visualized with 5 min color development, *HOXA5* was visualized at 25 min of color development, and an additional color intensification was also applied for *HOXA7* signal detection. For comparison, sections hybridized with a control sense RNA probe showed no signal (see Fig 3R). All the → point to developing hair follicles except the → in (L), which denotes the weak expression of *HOXA5* in the granular layer of the adult skin. Scale bar, 100 μ m.



in newborn and adult sebaceous glands, whereas expression of the other *HOX* genes was not detected (data not shown).

DISCUSSION

The *HOX* homeobox genes appear to be important developmental regulatory genes in animals as diverse as *Drosophila* to man (Krumlauf, 1992). These 39 genes encode transcription factors that appear to both activate and repress embryonic gene expression. Given these roles in other tissues, we and others have anticipated that the *HOX* genes might play roles in skin development (Scott and Goldsmith, 1993). Using a PCR-based screening method, the most commonly detected *HOX*-gene transcripts in human skin were *HOXA4*, *HOXA5*, and *HOXA7*. In addition, low level expression of several additional *HOXA* and *HOXB* genes as well as the *HOXC4* gene were detected by PCR in the fetal or adult skin samples. *In situ* hybridization analysis confirmed that *HOXA4* was expressed at high levels in developing skin. In addition, *HOXC4* was expressed at high levels in developing skin whereas *HOXA5*, *HOXA7*, and *HOXB7* were expressed at lower levels, and *HOXB4* was barely detected. Our observations confirm previous studies that have described detection of *HOXA7* (Thomas *et al*, 1989) and *HOXB7* (Simeone *et al*, 1987) and localization of *HOXC4* expression to the suprabasal layer of normal adult human skin (Reiger *et al*, 1994).

Taken together the two methods provide several important conclusions concerning expression of *HOX* genes during skin development. First, there appears to be a relatively conserved temporal and spatial expression pattern shared by many of the *HOX* genes. Expression appears to be limited to the basal cell layers in early skin development, prevalent in the basal and stratum granulosum layers of the epidermis as well as in the dermis during mid- to late second trimester develop-

ment, and dramatically downregulated and restricted to the granular layers of the adult epidermis. No *HOX* gene expression was detected in adult dermis by either RT-PCR or *in situ* hybridization analysis among the subset of *HOX* genes analyzed. On the other hand, *HOX* genes appear to be expressed in unique patterns in the developing hair follicles. Secondly, the relative levels expression for the various *HOX* genes appeared to be constant throughout development. The *in situ* analysis showed, for instance, that *HOXA4* was always detected at a relatively high level in every tissue and time point analyzed, whereas *HOXB4* was only weakly expressed in each specimen examined. In addition, the PCR analysis showed the three *HOXA* genes to be the predominant genes detected at all time points studied.

It is unclear what the function(s) of the Hox proteins might be during skin development. In other developmental programs, Hox proteins appear to specify positional and timing information (Krumlauf, 1994). There have been several reports of differential *HOX* gene expression patterns that support this type of role for Hox proteins in skin and hair development (Chuong *et al*, 1990; Bieberich *et al*, 1991; Kanzler *et al*, 1994); however, there is increasing evidence that enforced expression of these genes in other tissues can lead to unregulated proliferation (Aberdam *et al*, 1991; Maulbecker and Gruss, 1993; Thorsteinsdottir *et al*, 1997), whereas loss of *HOX* gene expression is associated with increased apoptosis.² In the context of skin development, *HOX* proteins might function as gene activators to upregulate cellular proliferation in the early developing epidermis. Similar proliferative

²Izon DJ, Komuves LG, Rozenfeld S, Fong S, Largman C, Lawrence HJ: Loss of function of the homeobox gene HoxA-9 retards T-cell development and induces cell death. Manuscript submitted.




Figure 3. Expression of *HOXB4*, *HOXB7*, and *HOXC4* in developing skin. *In situ* hybridization analysis was used to localize *HOX* homeobox gene expression during embryonic skin development. For parts (A)–(L), expressions in 10 wk (left panels), 17 wk (middle panels), and 21 wk (right panels) scalp skin are shown. *HOX* genes studied were: *HOXA4* for comparison and for visualization in the absence of hematoxylin counter-staining (A–C); *HOXB4* (D–F); *HOXB7* (G–I); and *HOXC4* (J–L). For parts (B)–(L), the → point to developing hair follicles. Expression of *HOX* genes in hair follicles is also shown in (M)–(Q), as follows: *HOXA4* expression (black →) in a hair follicle from 21 wk skin (M), the white → denotes the lack of expression in the dermal papilla; the parallel section hybridized with a sense control probe to show the background due to the developing hair shaft (→) (N); *HOXA4* expression in a hair follicle (black →) and sweat glands (white →) from newborn skin (O); *HOXA5* expression in the inner root sheath (→) of the hair follicle from newborn skin (P); *HOXC4* expression in the developing hair follicle (→) in 21 wk skin (Q). Part (R) was hybridized with a control sense *HOXA5* RNA probe. Sections (A)–(L) were adjacent to those stained with hematoxylin to visualize skin development shown in Fig 2(A,C,D), respectively. Parts (M) and (O) were counter-stained with hematoxylin to visualize morphologic structures. Expression of these genes was visualized identically using a standard 5 min color development. Scale bar, 100 μm.

functions might also be associated with the increased activity associated with the dermis during second trimester skin development. In this regard, it should be noted that first and early second trimester skin possesses the capacity to heal wounds without scar formation, whereas third trimester and adult skin lack this attribute (Weeks and Nath, 1993; Nath *et al*, 1995). It is possible that *HOX* proteins within the dermis provide information for proper regeneration of the wound during embryonic development. Finally, we hypothesize that *HOX* proteins might function as repressors to downregulate cellular proliferation in the outer layers of the mature skin. Although few biologic targets of *HOX* protein regulation have been identified, several of the suggested targets of homeodomain protein action are cellular adhesion or signaling molecules, which could play a role(s) in cellular proliferation (Jones *et al*, 1992; White *et al*, 1992; Tomotsune *et al*, 1993).

The hypothetical roles described above would require *HOX* proteins to act as activators in one cellular environment and as repressors in a different cellular layer. One mechanism by which the *HOX* transcription factors might exhibit different phenotypic effects is through the presence or absence of partner proteins. We and others have recently shown that *HOX* proteins form cooperative DNA binding complexes with members of the PBX family of non-*HOX* homeodomain proteins (Chang *et al*, 1995, 1996; Lu *et al*, 1995; Phelan *et al*, 1995; van Dijk *et al*, 1995). Furthermore, a recent model for *HOX* protein function proposes that they cooperatively bind DNA as activators in the presence of PBX and function as transcriptional repressors by binding alone to DNA (Pinsonneault *et al*, 1997). Thus one model would be that the *HOX* proteins might function as activators in early skin development by binding DNA with PBX or other protein partners, and might

function as repressors in the granular cells of the adult epidermis if the array of partner proteins was different in these cells.

It is somewhat puzzling that there appears to be such a high redundancy of *HOX* gene expression in the developing skin. While *Hox* genes are expressed in developing murine spinal cord or limb buds in redundant, overlapping patterns, there are also apparent areas in which individual *Hox* genes show increased relative expression (Krumlauf, 1994). This does not appear to be the case in the developing skin, because the expression patterns for the six *HOX* genes examined appeared very similar. In addition, the expression of each gene appeared to be relatively uniform across the body. Although the level of our analysis was insufficient to determine if multiple *HOX* genes were expressed in individual cells, it appears likely that there is a high level of redundancy at the cellular level. Although this complexity is difficult to interpret in terms of functional roles for the homeoproteins, it should be noted that signaling molecules such as BMP-2 and BMP-4 (Francis *et al*, 1994), or the Wnt-like proteins (Parr and McMahon, 1994) that are thought to be both regulators and perhaps targets of *HOX* proteins (Iler *et al*, 1995; Manak *et al*, 1995), are themselves expressed in complex patterns throughout vertebrate development. In addition, functional *Hox* gene redundancy in murine rhombomere formation has recently been demonstrated (Gould *et al*, 1997). While the purpose of this redundancy is not clear, we have recently demonstrated large differences in complex stabilities between various *HOX* proteins bound to DNA with PBX (Shen *et al*, 1997). We hypothesize that the protein products derived from the *HOX* genes, which we have demonstrated to be expressed in developing skin, function by competing for DNA binding sites both as complexes with a variable

set of partner proteins and as independent DNA binding proteins. As a first step in defining the regulatory targets for HOX proteins in the skin, it will be necessary to elucidate the set of DNA binding partners expressed throughout skin development.

This work was supported by a grant from the American Plastic Surgery Education Foundation/Ethicon Research Fellowship to Eric Stelnicki, by the Research Service of the Department of Veterans Affairs (C.L.), NIH Grant #HD-25505 (S.A.), and by the G. Harold and Leila Y. Mathers Charitable Foundation (M.H.). We wish to thank members of the Largman laboratory for assistance and helpful comments. and Dr. Peter Elias for reading the manuscript. We also wish to thank the following individuals for supplying human HOX cDNA clones: Dr. Judy Gasson (HOXA5), Dr. Debbie Wolgemuth (HOXA4), and Dr. Eduardo Boncinelli (HOXA7, HOXB4, and HOXC4). A partial genomic clone containing the 3' nonhomeobox coding and untranslated regions of human HOXA7 was the kind gift of Dr. Guy Sauvageau.

REFERENCES

- Aberdam D, Negreanu V, Sachs L, Blatt C: The oncogenic potential of an activated Hox-2.4 homeobox gene in mouse fibroblasts. *Mol Cell Biol* 11:554-557, 1991
- Bieberich CJ, Ruddle FH, Stenn KS: Differential expression of the Hox 3.1 gene in adult mouse skin. *Ann N Y Acad Sci* 642:346-354, 1991
- Chang C-P, Shen W-F, Rozenfeld S, Lawrence HJ, Largman C, Cleary ML: Pbx proteins display hexapeptide-dependent cooperative DNA binding with a subset of Hox proteins. *Genes Dev* 9:663-674, 1995
- Chang C-P, Brocchieri L, Shen W-F, Largman C, Cleary ML: Pbx modulation of Hox homeodomain N-terminal arms establishes a gradient of DNA-binding specificities across the Hox locus. *Mol Cell Biol* 16:1734-1745, 1996
- Chomczynski P, Sacchi N: Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 162:156-159, 1987
- Chuong CM, Oliver G, Ting SA, Jegalian BG, Chen HM, De Robertis EM: Gradients of homeoproteins in developing feather buds. *Development* 110:1021-1030, 1990
- Detmer K, Lawrence HJ, Largman C: Expression of class I homeobox genes in fetal and adult murine skin. *J Invest Dermatol* 101:517-522, 1993
- van Dijk MA, Peltenburg LTC, Murre C: Hox gene products modulate the DNA binding activity of Pbx1 and Pbx2. *Mech Dev* 52:99-108, 1995
- Francis PH, Richardson MK, Brickell PM, Tickle C: Bone morphogenetic proteins and a signalling pathway that controls patterning in the developing chick limb. *Dev* 120:209-218, 1994
- Gehring WJ, Affolter M, Burglin T: Homeodomain proteins. *Ann Rev Biochem* 63:487-526, 1994
- van Genderen C, Okamura RM, Farinas I, Quo R-G, Parslow TG, Bruhn L, Grosschedl R: Development of several organs that require inductive epithelial-mesenchymal interactions is impaired in LEF-1 deficient mice. *Genes Dev* 8:2691-2703, 1994
- Gould A, Morrison A, Sproat G, White RAH, Krumlauf R: Positive cross-regulation and enhancer sharing: two mechanisms for specifying overlapping Hox expression patterns. *Genes Dev* 11:900-913, 1997
- Holleran WM, Takagi Y, Imokawa G, Jackson S, Lee JM, Elias PM: beta-Glucocerebrosidase activity in murine epidermis: characterization and localization in relation to differentiation. *J Lipid Res* 33:1201-1209, 1992
- Iler N, Rowitch DH, Echelard Y, McMahon AP, Abate-Shen C: A single homeodomain binding site restricts spatial expression of Wnt-1 in the developing brain. *Mech Dev* 53:87-96, 1995
- Jones FS, Chalepakos G, Gruss P, Edelman GM: Activation of the cytotoxic promoter by the homeobox-containing gene Evx-1. *Proc Natl Acad Sci USA* 89:2091-2095, 1992
- Kanzler B, Viallet JP, Le Mouellic H, Boncinelli E, Duboule D, Dhoubailly D: Differential expression of two different homeobox gene families during mouse tegument morphogenesis. *Int J Dev Biol* 38:633-640, 1994
- Krumlauf R: Evolution of the vertebrate Hox homeobox genes. *Bioessays* 14:245-252, 1992
- Krumlauf R: Hox genes in vertebrate development. *Cell* 78:191-201, 1994
- Lu Q, Knoepfner PS, Scheele J, Wright DD, Kamps MP: Both Pbx1 and E2A-Pbx1 bind the DNA motif ATCAATCAA cooperatively with the products of multiple murine Hox genes, some of which are themselves oncogenes. *Mol Cell Biol* 15:3786-3795, 1995
- Luetke NC, Qiu TH, Peiffer RL, Oliver P, Smithes O, Lee DC: TGF α deficiency results in hair follicle and eye abnormalities in targeted and waved-1 mice. *Cell* 73:263-278, 1993
- Manak JR, Mathies LD, Scott MP: Regulation of a decapentaplegic midgut enhancer by homeotic proteins. *Dev* 120:3605-3619, 1995
- Mann GB, Fowler KJ, Anastasia G, Nice EC, Williams RL, Dunn AR: Mice with a null mutation in the TGF α gene have abnormal skin architecture, wavy hair, and curly whiskers and often develop corneal inflammation. *Cell* 73:249-261, 1993
- Mathews CH, Detmer K, Lawrence HJ, Largman C: Expression of the Hox 2.2 homeobox gene in murine embryonic epidermis. *Differentiation* 52:177-184, 1993
- Maulbecker CC, Gruss P: The oncogenic potential of deregulated homeobox genes. *Cell Growth Differen* 4:431-441, 1993
- Miettinen PJ, Berger JE, Meneses J, Phung Y, Pedersen RA, Werb Z, Derynck R: Epithelial immaturity and multiorgan failure in mice lacking epidermal growth factor receptor. *Nature* 376:337-341, 1995
- Murillas R, Larcher F, Conti CJ, Santos M, Ulrich A, Jorcano JL: Expression of a dominant negative mutant of epidermal growth factor in the epidermis of transgenic mice elicits striking alterations in hair follicle development and skin structure. *EMBO J* 14:5216-5223, 1995
- Nath RK, Jensen J, Weeks PM: Temporally variable induction of homeobox gene D 11 during wound repair in the fetal mouse limb. *Surgical Forum: American College of Surgeons* XL:VI, 1995
- Nohno T, Ishikawa T, Saito T, Hosokawa K, Noji S, Wolsing DH, Rosenbaum JS: Identification of a human type II receptor for bone morphogenetic protein-4 that forms differential heteromeric complexes with bone morphogenetic protein type I receptors. *J Biol Chem* 270:22522-22526, 1995a
- Nohno T, Kawakami Y, Ohuchi H, Fujiwara A, Yoshioka H, Noji S: Involvement of the Sonic hedgehog gene in chick feather formation. *Biochem Biophys Res Commun* 206:33-39, 1995b
- Parr BA, McMahon AP: Wnt genes and vertebrate development. *Curr Opin Genet Dev* 4:523-528, 1994
- Phelan ML, Rambaldi I, Featherstone MS: Cooperative interactions between HOX and PBX proteins mediated by a conserved peptide motif. *Mol Cell Biol* 15:3989-3997, 1995
- Pinsonneault J, Florence B, Vaessin H, McGinnis W: A model for extradenticle function as a switch that changes HOX proteins from repressors to activators. *EMBO J* 16:2032-2042, 1997
- Reiger E, Bijl JJ, van Oostveen JW, et al: Expression of the homeobox gene HOXC4 in keratinocytes of normal skin and epithelial skin tumors is correlated with differentiation. *J Invest Dermatol* 103:341-346, 1994
- Sauvageau G, Lansdorp PM, Eaves CE, et al: Differential expression of homeobox genes in functionally distinct CD34+ subpopulations of human bone marrow cells. *Proc Natl Acad Sci USA* 91:12223-12227, 1994
- Scott GA, Goldsmith LA: Homeobox genes and skin development: a review. *J Invest Dermatol* 101:3-8, 1993
- Shen W-F, Rozenfeld S, Lawrence HJ, Largman C: The Abd-B-like Hox homeodomain proteins can be subdivided by the ability to form complexes with Pbx1a on a novel DNA target. *J Biol Chem* 272:8198-8206, 1997
- Simeone A, Mavilio F, Acampora D, et al: Two human homeobox genes, C1 and C8: Structure analysis and expression in embryonic development. *Proc Natl Acad Sci USA* 84:4914-4918, 1987
- Thomas NE, Polakowska R, Rogers KE, Kelly E, Goldsmith LA: Homeobox containing genes found in an adult human keratinocyte library. *Clin Res* 37:595A, 1989
- Thorsteinsdottir U, Sauvageau G, Hough MR, et al: Overexpression of HOXA10 in murine hematopoietic cells perturbs both myeloid and lymphoid differentiation and leads to acute myeloid leukemia. *Mol Cell Biol* 17:495-505, 1997
- Tomotsune D, Shoji H, Wakamatsu Y, Kondoh H, Takahashi N: A mouse homologue of the Drosophila tumour-suppressor gene I (2) gl controlled by Hox-C8 *in vivo*. *Nature* 365:69-72, 1993
- Weeks PM, Nath RK: Fetal wound repair: a new direction. *Plast Reconstr Surg* 91:922-924, 1993
- White RA, Brookman JJ, Gould AP, Meadows LA, Sahahidhara LS, Strutt DI, Weaver TA: Targets of homeotic gene regulation in Drosophila. *J Cell Sci Suppl* 16:53-60, 1992
- Wilkinson DG: In situ hybridization. In: Rickwood D, Hames BD (eds). *The Practical Approach Series*. IRL Press, London, 1992
- Zeltser L, Desplan C, Heintz N: Hoxb-13: a new Hox gene in a distant region of the HOXB cluster maintains colinearity. *Dev* 122:2475-2484, 1996
- Zhou P, Burne C, Jacobs J, Fuchs E: Lymphoid enhancer factor 1 directs hair follicle patterning and epithelial cell fate. *Genes Dev* 9:700-713, 1995