

**NEWS RELEASE  
FOR IMMEDIATE RELEASE**

**Media contact:**

Theresa Monturano, Senior Publisher  
Elsevier  
+1 215 239 3711  
[hmsmedia@elsevier.com](mailto:hmsmedia@elsevier.com)

**Increased depression, suicidal thoughts and stress are reported in patients with chronically itchy skin**

*A new multicenter European study that assessed the burden of itch on dermatological patients' psychological wellbeing is reported in the Journal of Investigative Dermatology*

**Philadelphia, October 29, 2019** – Itch is a very common symptom in patients suffering from skin diseases. In a new multicenter cross-sectional [study](#) on the psychological burden of itch in the [Journal of Investigative Dermatology](#), published by Elsevier, investigators report that the presence of itch in dermatological patients was significantly associated with clinical depression, suicidal ideation and stress. They recommend providing patients with access to a multidisciplinary team to prevent and manage problems associated with itch.

The burden of itch has already been described in conjunction with a number of specific skin diseases including hand eczema; psoriasis; nodular prurigo (a skin disease that causes hard, itchy lumps to form on the skin); hidradenitis suppurativa (a painful, long-term skin condition that causes abscesses and scarring on the skin) among hemodialysis patients; and in chronic itch patients in general.

“There are already studies showing evidence of a correlation between itch and mental health problems in general, and in specific skin disorders, but there is a lack of a cross-sectional study across chronic skin diseases,” explained lead investigator Florence J. Dalgard, MD, PhD, Department of Dermatology and Venereology, Skåne University Hospital, Lund University, Malmö, Sweden.

Part of a large European multicenter study conducted by the European Society for Dermatology and Psychiatry (ESDaP), the present study compared the psychological burden of disease and health-related quality of life between dermatological patients with itch and those without itch, as well as with healthy controls.

Investigators collected data from dermatological clinics in 13 European countries on 3,530 patients with skin diseases and compared the results with more than 1,000 healthy controls. Patients completed

questionnaires and were also examined clinically. Outcome measures included the presence, chronicity and intensity of itch; the Hospital Anxiety and Depression Scale; sociodemographics; suicidal ideation, and stress, including negative life events; and economic difficulties.

The prevalence of itch in dermatological conditions was nearly 90 percent in prurigo and related conditions; 86 percent in atopic dermatitis; 82 percent in hand eczema; 78 percent in other eczema; 76 percent in urticarial; and 70 percent in psoriasis.



Caption: Scratching that itch. Itchy patients were more likely to be seriously depressed and to have suicidal thoughts.

The prevalence of depression was 14 percent in patients with itch compared to 5.7 percent in patients without itch, six percent in controls with itch, and three percent in controls without itch. The prevalence of suicidal thoughts was 15.7 percent in patients with itch, nine percent in patients without itch, 18.6 percent in controls with itch and 8.6 percent in controls without itch. The reported occurrence of stressful life events was higher in individuals with itch than in those without itch. Patients with itch were also likely to experience more economic problems.

“Our research shows that itch has a high impact on quality of life,” commented Dr. Dalgard. “This study illustrates the burden of the symptom of itch and its multidimensional aspect. The management of patients with itch should involve access to a multidisciplinary team when necessary, as is already the case in several European countries.”

The investigators also recommend preventive measures, such as psoriasis education programs or targeted web-based information. In many chronic inflammatory skin disorders, early aggressive treatment tailored specifically for the patient might help to reduce itch at the earliest possible opportunity and prevent the development of mental health problems. Existing anti-itch interventions should be implemented more frequently in the routine care of dermatological patients.

---

#### **Notes for editors**

The article is “Itch and mental health in dermatological patients across Europe: a cross sectional study in 13 countries,” by Florence J. Dalgard, Åke Svensson, Jon Anders Halvorsen , Uwe Gieler, Christina Schut, Lucia Tomas-Aragones, Lars Lien, Françoise Poot, Gregor B. E. Jemec, Laurent Misery, Csanad Szabo, Dennis Linder, Francesca Sampogna, Saskia Spillekom-van Kouli, Flora Balieva, Jacek C.

Szepietowski, Andrey Lvov, Servando E. Marron, Ilknur K. Altunay, Andrew Y. Finlay, Sam Salek and Jörg Kupfer(<https://doi.org/10.1016/j.jid.2019.05.034>). It appears in the *Journal of Investigative Dermatology* published by Elsevier. The article is openly available at [https://www.jidonline.org/article/S0022-202X\(19\)33209-9/fulltext](https://www.jidonline.org/article/S0022-202X(19)33209-9/fulltext).

This study was supported by South-Eastern Norway Regional Health Authority in Norway, Skåne University Hospital, Malmö in Sweden and Justus Liebig University in Giessen, Germany.

Full text of this article is available to credentialed journalists upon request; contact Theresa Monturano, Senior Publisher, Elsevier, at +1 215 239 3711 or [hmsmedia@elsevier.com](mailto:hmsmedia@elsevier.com). Journalists wishing to interview the authors should contact Florence J. Dalgard at [florikje@gmail.com](mailto:florikje@gmail.com).

### **About *Journal of Investigative Dermatology***

The *Journal of Investigative Dermatology* (JID) publishes high impact reports describing original research related to all aspects of cutaneous biology and skin disease. Descriptions of important findings that result from basic, translational, or clinical research are published. Clinical research can include, but is not limited to, interventional trials, genetics studies, epidemiology, and health services research. [www.jidonline.org](http://www.jidonline.org)

### **About Elsevier**

Elsevier is a global information analytics business that helps scientists and clinicians to find new answers, reshape human knowledge, and tackle the most urgent human crises. For 140 years, we have partnered with the research world to curate and verify scientific knowledge. Today, we're committed to bringing that rigor to a new generation of platforms. Elsevier provides digital solutions and tools in the areas of strategic research management, R&D performance, clinical decision support, and professional education; including ScienceDirect, Scopus, SciVal, ClinicalKey and Sherpath. Elsevier publishes over 2,500 digitized journals, including *The Lancet* and *Cell*, 39,000 e-book titles and many iconic reference works, including *Gray's Anatomy*. Elsevier is part of RELX, a global provider of information-based analytics and decision tools for professional and business customers. [www.elsevier.com](http://www.elsevier.com)